

Parallel Scientific Computing

Course AMS301 — Fall 2022 — Lecture 6

Iterative methods for linear systems (2)
Nonstationary iterative methods

Find $\mathbf{x} \in \mathbb{R}^n$ such that $\boxed{\mathbf{Ax} = \mathbf{b}}$ with $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{b} \in \mathbb{R}^n$.

Solution procedures

- ▶ **Direct methods:** Factorization of \mathbf{A} into triangular and diagonal matrices (ex. $\mathbf{A} = \mathbf{LU}$) and solution of simpler problems.

$$\mathbf{Ax} = \mathbf{b} \quad \Leftrightarrow \quad \mathbf{LUx} = \mathbf{b} \quad \Leftrightarrow \quad \begin{cases} \mathbf{Ly} = \mathbf{b} \\ \mathbf{Ux} = \mathbf{y} \end{cases}$$

Advantages: exact solution known after a given number of operations

Difficulties: heavy computational cost (*operations/memory*), hard to parallelize

- ▶ **Iterative methods:** Iterative procedure to minimizing an error $\|\mathbf{x}^{(\ell)} - \mathbf{x}_{\text{ref}}\|$ and/or a residual $\|\mathbf{Ax}^{(\ell)} - \mathbf{b}\|$.

$$\begin{cases} \mathbf{x}^{(0)} = \text{Iter}_{(0)}(\mathbf{A}, \mathbf{b}) \\ \mathbf{x}^{(\ell+1)} = \text{Iter}^{(\ell+1)}(\mathbf{x}^{(\ell)}, \mathbf{x}^{(\ell-1)}, \dots, \mathbf{A}, \mathbf{b}), \quad \text{pour } \ell \geq 0 \end{cases}$$

Advantages: limited cost per iteration (*operations/memory*), easy to parallelize

Difficulties: approximate solution, control of the convergence of the process

Find $\mathbf{x} \in \mathbb{R}^n$ such that $\boxed{\mathbf{Ax} = \mathbf{b}}$ with $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{b} \in \mathbb{R}^n$.

► **Iterative methods:**

$$\left| \begin{array}{l} \mathbf{x}^{(0)} = \text{Iter}_{(0)}(\mathbf{A}, \mathbf{b}) \\ \mathbf{x}^{(\ell+1)} = \text{Iter}^{(\ell+1)}(\mathbf{x}^{(\ell)}, \mathbf{x}^{(\ell-1)}, \dots, \mathbf{A}, \mathbf{b}), \quad \text{pour } \ell \geq 0 \end{array} \right.$$

The **order** of the method is the numb. of steps which the current iter. depends on.
Stationary method if the functions $\text{Iter}^{(\ell)}$ are indep. of ℓ , otherwise **nonstationary**
Linear method if the functions $\text{Iter}^{(\ell)}$ are linear, otherwise **nonlinear**

In a previous session, we considered statio. linear iterative schemes of first order:

$$\left| \begin{array}{l} \mathbf{x}^{(0)} \text{ given} \\ \mathbf{x}^{(\ell+1)} = \mathbf{B}\mathbf{x}^{(\ell)} + \mathbf{f}, \quad \ell \geq 0 \end{array} \right.$$

where $\mathbf{B} \in \mathbb{R}^{n \times n}$ is the **iteration matrix** and $\mathbf{f} \in \mathbb{R}^n$ depends on \mathbf{b} .

Stationary linear iterative method of first order

Regular decomposition: $\mathbf{A} = \mathbf{M} - \mathbf{N}$ where $\mathbf{M} \in \mathbb{R}^{n \times n}$ is invertible.

Stationary method	
$\mathbf{x}^{(0)} \in \mathbb{C}^n$ for $\ell = 0, 1, \dots$ do $\mathbf{M}\mathbf{x}^{(\ell+1)} = \mathbf{N}\mathbf{x}^{(\ell)} + \mathbf{b}$ end	i.e. $\mathbf{x}^{(\ell+1)} = \mathbf{B}\mathbf{x}^{(\ell)} + \mathbf{f}$ with $\mathbf{B} = \mathbf{M}^{-1}\mathbf{N}$ and $\mathbf{f} = \mathbf{M}^{-1}\mathbf{b}$

Choices

	By points	By blocks
Jacobi	$\mathbf{M} = \mathbf{D}$	$\mathbf{M} = \mathbf{D}^{\text{blk}}$
Gauss-Seidel	$\mathbf{M} = \mathbf{D} + \mathbf{L}$	$\mathbf{M} = \mathbf{D}^{\text{blk}} + \mathbf{L}^{\text{blk}}$

$$\underbrace{\begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}}_{\mathbf{A}} = \underbrace{\begin{bmatrix} \times & & \\ & \times & \\ & & \times \end{bmatrix}}_{\mathbf{D}} + \underbrace{\begin{bmatrix} & \times & \\ \times & \times & \\ & & \end{bmatrix}}_{\mathbf{L}} + \underbrace{\begin{bmatrix} & \times & \times \\ & & \times \\ & & \end{bmatrix}}_{\mathbf{U}}$$

Solution of linear systems — Nonstationary methods

Find $\mathbf{x} \in \mathbb{R}^n$ such that $\boxed{\mathbf{Ax} = \mathbf{b}}$ with $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{b} \in \mathbb{R}^n$.

In a previous session, we considered statio. linear iterative schemes of first order:

$$\left| \begin{array}{l} \mathbf{x}^{(0)} \text{ given} \\ \mathbf{x}^{(\ell+1)} = \mathbf{B}\mathbf{x}^{(\ell)} + \mathbf{f}, \quad \ell \geq 0 \end{array} \right.$$

where $\mathbf{B} \in \mathbb{R}^{n \times n}$ is the **iteration matrix** and $\mathbf{f} \in \mathbb{R}^n$ depends on \mathbf{b} .

During this session, we consider **nonstationary linear iterative schemes** of the form:

$$\left| \begin{array}{l} \mathbf{x}^{(0)} \text{ given} \\ \mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)}, \quad \ell \geq 0 \end{array} \right.$$

with the **step** $\alpha^{(\ell)}$ and the **direction** $\mathbf{p}^{(\ell)}$ must be chosen.

Nonstationary iterative methods for linear systems

Conjugate gradient method

Interlude on Krylov spaces

Few words about GMRES

Find $\mathbf{x} \in \mathbb{R}^n$ such that $\mathbf{Ax} = \mathbf{b}$ with $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{b} \in \mathbb{R}^n$,
where \mathbf{A} is symmetric positive definite (SPD).

$$\mathbf{A} = \mathbf{A}^\top \quad \text{and} \quad (\mathbf{A}\mathbf{v}, \mathbf{v}) > 0, \quad \forall \mathbf{v} \in \mathbb{R}^n \setminus \{0\}$$

Link with a minimization problem

We consider the following minimization problem:

Find $\mathbf{x} \in \mathbb{R}^n$ that minimizes the functional $J(\mathbf{v}) = \frac{1}{2}(\mathbf{A}\mathbf{v}, \mathbf{v}) - (\mathbf{b}, \mathbf{v})$.

If \mathbf{A} is an SPD matrix:

- ▶ The functional $J(\mathbf{v})$ is strictly convex on \mathbb{R}^n .
- ▶ The functional $J(\mathbf{v})$ has a unique minimum.
- ▶ The minimum of $J(\mathbf{v})$, denoted \mathbf{v}_{\min} , is such that $\nabla J|_{\mathbf{v}_{\min}} = 0$ and $\mathbf{A}\mathbf{v}_{\min} = \mathbf{b}$.


Solving the minimization problem is equivalent to solving the system!

Conjugate gradient method — Principle [2/2]

Principle of the conjugate gradient method

- ▶ Starting from a vector $\mathbf{x}^{(0)}$, we compute vectors $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$, ... such that J is min.
- ▶ At each iteration, we take one step $\alpha^{(\ell)}$ along direction $\mathbf{p}^{(\ell)}$:

$$\mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)} = \mathbf{x}^{(0)} + \sum_{i=0}^{\ell} \alpha^{(i)} \mathbf{p}^{(i)}$$



Source: Wikipedia

Steepest descent method

At each step:

- ▶ Choice of direction $\mathbf{p}^{(\ell)}$ \rightarrow We take the gradient: $\mathbf{p}^{(\ell)} = -\nabla J(\mathbf{x}^{(\ell)})$
- ▶ Choice of step $\alpha^{(\ell)}$ \rightarrow We take the one that minimizes $J(\mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)})$.

We have

$$\left\{ \begin{array}{l} J(\mathbf{v}) = \frac{1}{2} (\mathbf{A}\mathbf{v}, \mathbf{v}) - (\mathbf{b}, \mathbf{v}) \\ \nabla J(\mathbf{v}) = \mathbf{A}\mathbf{v} - \mathbf{b} \quad (\text{Then, } \mathbf{p}^{(\ell)} = \mathbf{b} - \mathbf{A}\mathbf{x}^{(\ell)}, \text{ which is the residual!}) \\ \min_{\alpha} J(\mathbf{x}^{(\ell)} + \alpha \mathbf{p}^{(\ell)}) \Leftrightarrow \alpha = (\mathbf{b} - \mathbf{A}\mathbf{x}^{(\ell)}, \mathbf{p}^{(\ell)}) / (\mathbf{A}\mathbf{p}^{(\ell)}, \mathbf{p}^{(\ell)}) \end{array} \right.$$

Steepest descent method

$$\mathbf{x}^{(0)} \in \mathbb{R}^n$$

$$\mathbf{p}^{(0)} = \mathbf{b} - \mathbf{A}\mathbf{x}^{(0)}$$

for $\ell = 0, 1, \dots$ **do**

$$\alpha^{(\ell)} = (\mathbf{p}^{(\ell)}, \mathbf{p}^{(\ell)}) / (\mathbf{A}\mathbf{p}^{(\ell)}, \mathbf{p}^{(\ell)})$$

Comput. of step

$$\mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)}$$

Update

$$\mathbf{p}^{(\ell+1)} = \mathbf{b} - \mathbf{A}\mathbf{x}^{(\ell+1)}$$

Comput. of direction/residual

if $\|\mathbf{p}^{(\ell+1)}\| \leq \varepsilon \|\mathbf{p}^{(0)}\|$ **then break**

end

Conjugate gradient method

- ▶ We take $\{\mathbf{p}^{(\ell)}\}_{\ell=0\dots n-1}$ such that they form a basis of \mathbb{R}^n .
We take a **A-orthogonal** basis, i.e. orthogonal with the scalar product $(\mathbf{A}\cdot, \cdot)$:

$$(\mathbf{A}\mathbf{p}^{(i)}, \mathbf{p}^{(j)}) = 0, \quad \forall i \neq j$$

- ▶ We take $\{\alpha^{(\ell)}\}_{\ell=0\dots n-1}$ such that

$$\mathbf{x} = \mathbf{x}^{(0)} + \sum_{i=0}^{n-1} \alpha^{(i)} \mathbf{p}^{(i)}$$

where \mathbf{x} is the solution of the problem.

Conjugate gradient method

```

 $\mathbf{x}^{(0)} \in \mathbb{R}^n$ 
 $\mathbf{p}^{(0)} = \mathbf{b} - \mathbf{A}\mathbf{x}^{(0)}$ 
for  $\ell = 0, 1, \dots$  do
  |
  |  $\vdots$ 
  |  $\mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)} \quad = \mathbf{x}^{(0)} + \sum_{i=0}^{\ell} \alpha^{(i)} \mathbf{p}^{(i)}$ 
  |  $\vdots$ 
end
  Update
  
```

Conjugate gradient method (*continuation*)

At each step:

- ▶ Choice of direction $\mathbf{p}^{(\ell)}$ \rightarrow Part of $\mathbf{r}^{(\ell)}$ that is \mathbf{A} -orthogonal with $\mathbf{p}^{(\ell-1)}$
- ▶ Choice of step $\alpha^{(\ell)}$ \rightarrow Value that minimizes $J(\mathbf{x}^{(\ell)} + \alpha^{(\ell)}\mathbf{p}^{(\ell)})$

Conjugate gradient method		
$\mathbf{x}^{(0)} \in \mathbb{R}^n$		
$\mathbf{r}^{(0)} = \mathbf{b} - \mathbf{A}\mathbf{x}^{(0)}$		
$\mathbf{p}^{(0)} = \mathbf{r}^{(0)}$		
for $\ell = 0, 1, \dots$ do		
$\alpha^{(\ell)} = (\mathbf{r}^{(\ell)}, \mathbf{p}^{(\ell)}) / (\mathbf{A}\mathbf{p}^{(\ell)}, \mathbf{p}^{(\ell)})$		Comput. of step
$\mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)}\mathbf{p}^{(\ell)}$	$= \mathbf{x}^{(0)} + \sum_{i=0}^{\ell} \alpha^{(i)}\mathbf{p}^{(i)}$	Update
$\mathbf{r}^{(\ell+1)} = \mathbf{r}^{(\ell)} - \alpha^{(\ell)}\mathbf{A}\mathbf{p}^{(\ell)}$	$= \mathbf{b} - \mathbf{A}\mathbf{x}^{(\ell+1)}$	Comput. of residual
$\beta^{(\ell)} = -(\mathbf{A}\mathbf{r}^{(\ell+1)}, \mathbf{p}^{(\ell)}) / (\mathbf{A}\mathbf{p}^{(\ell)}, \mathbf{p}^{(\ell)})$		
$\mathbf{p}^{(\ell+1)} = \mathbf{r}^{(\ell+1)} + \beta^{(\ell)}\mathbf{p}^{(\ell)}$		Comput. of direction
if $\ \mathbf{r}^{(\ell+1)}\ \leq \varepsilon \ \mathbf{r}^{(0)}\ $ then break		
end		

Theoretical aspects

- ▶ Method for symmetric pos. def. (SDP) and Hermitian pos. def. (HPD) matrices
- ▶ All the directions are \mathbf{A} -orthogonal.
- ▶ By construction, **convergence with maximum n iterations!** (if ∞ accuracy)
- ▶ The error $\mathbf{e}^{(\ell)} = \mathbf{x} - \mathbf{x}^{(\ell)}$ verifies

$$\sqrt{(\mathbf{A}\mathbf{e}^{(\ell)}, \mathbf{e}^{(\ell)})} \leq \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^\ell \sqrt{(\mathbf{A}\mathbf{e}^{(0)}, \mathbf{e}^{(0)})}$$

with the **condition number** $\kappa = \|\mathbf{A}\| \|\mathbf{A}^{-1}\|$.

Because \mathbf{A} is SPD: $\kappa = \lambda_{\max}/\lambda_{\min}$. The max/min eigenvalues influence the convergence rate. When κ is close to 1, convergence is fast.

Algorithmic aspects

- ▶ Linear algebraic operations (BLAS 1 and 2) \Rightarrow Easy for parallel computing
- ▶ Computation of scalar products and norms \Rightarrow Collective communications

Best method for systems with SPD matrices!
Extensions for general matrices?

Nonstationary iterative methods for linear systems

Conjugate gradient method

Interlude on Krylov spaces

Few words about GMRES

Krylov spaces — Motivation

The conjugate gradient method relies on **adding an update term** to the current solution:

$$\mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)},$$

with the update direction $\mathbf{p}^{(\ell)}$ and the step $\alpha^{(\ell)}$.

The update term can be rewritten as:

$$\mathbf{x}^{(\ell+1)} - \mathbf{x}^{(0)} = \sum_{i=0}^{\ell} \alpha^{(i)} \mathbf{p}^{(i)}$$

It belongs to the subspace:

$$\text{span}(\mathbf{p}^{(0)}, \dots, \mathbf{p}^{(\ell)}) \subseteq \mathbb{R}^n$$

Toward general iterative methods ...

General iterative methods for more general non-symmetric/non-Hermitian matrices can be built by considering **Krylov subspaces**, e.g.

$$\mathcal{K}_{\ell}(\mathbf{A}, \mathbf{r}^{(0)}) := \text{span}(\mathbf{r}^{(0)}, \mathbf{A}\mathbf{r}^{(0)}, \dots, \mathbf{A}^{\ell-1}\mathbf{r}^{(0)}).$$

These methods rely on 2 steps:

1. Building a basis $\{\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(\ell)}\}$ for a Krylov subspace
2. Solving a minimization problem to get the update term towards the “*best solution*”

Definition – Krylov subspace

The **order- ℓ Krylov subspace associated to $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\mathbf{v} \in \mathbb{R}^n$** , with $\ell < n$, is the linear subspace spanned by the images of \mathbf{v} under the first ℓ powers of \mathbf{A} , starting from \mathbf{A}^0 :

$$\mathcal{K}_\ell(\mathbf{A}, \mathbf{v}) := \text{span}(\mathbf{v}, \mathbf{A}\mathbf{v}, \dots, \mathbf{A}^{\ell-1}\mathbf{v}).$$

Properties

- $\mathcal{K}_\ell(\mathbf{A}, \mathbf{v}) \subseteq \mathcal{K}_{\ell+i}(\mathbf{A}, \mathbf{v}) \subseteq \mathbb{R}^n, \quad \forall i \geq 0$
- $\mathbf{A}\mathcal{K}_\ell(\mathbf{A}, \mathbf{v}) \subseteq \mathcal{K}_{\ell+1}(\mathbf{A}, \mathbf{v}), \quad \forall \ell$
- $\dim(\mathcal{K}_\ell(\mathbf{A}, \mathbf{v})) = \min(\ell, \min. \text{ degree of non-zero poly. } \mathcal{P} \text{ such that } \mathcal{P}(\mathbf{A})\mathbf{v} = 0)$
- The sequence $(\mathcal{K}_\ell(\mathbf{A}, \mathbf{v}))_\ell$ is strictly increasing from 1 to ℓ_{\max} , then it is constant starting from ℓ_{\max} , where $\ell_{\max} := \operatorname{argmax}_\ell(\dim \mathcal{K}_\ell(\mathbf{A}, \mathbf{v}))$.

Example

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 4 & -2 \\ 2 & 2 & -1 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad \mathbf{A}\mathbf{v} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} \quad \mathbf{A}^2\mathbf{v} = \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathcal{K}_1(\mathbf{A}, \mathbf{v}) = \text{span}(\mathbf{v}) \quad \mathcal{K}_2(\mathbf{A}, \mathbf{v}) = \text{span}(\mathbf{v}, \mathbf{A}\mathbf{v}) \quad \mathcal{K}_3(\mathbf{A}, \mathbf{v}) = \text{span}(\mathbf{v}, \mathbf{A}\mathbf{v}, \mathbf{A}^2\mathbf{v})$$

$$\dim(\mathcal{K}_1(\mathbf{A}, \mathbf{v})) = 1 \quad \dim(\mathcal{K}_2(\mathbf{A}, \mathbf{v})) = 2 \quad \dim(\mathcal{K}_3(\mathbf{A}, \mathbf{v})) = 2$$

Recap:

$$\mathbf{x}^{(\ell+1)} = \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)}$$

$$\mathbf{r}^{(\ell+1)} = \mathbf{r}^{(\ell)} - \alpha^{(\ell)} \mathbf{A} \mathbf{p}^{(\ell)}$$

$$\mathbf{p}^{(\ell+1)} = \mathbf{r}^{(\ell+1)} + \beta^{(\ell)} \mathbf{p}^{(\ell)}$$

Property: $\mathbf{x}^{(\ell+1)} - \mathbf{x}^{(0)} \in \mathcal{K}_{\ell+1}(\mathbf{A}, \mathbf{r}^{(0)})$

Proof (for info)

One has:

$$\begin{aligned} \mathbf{x}^{(\ell+1)} &= \mathbf{x}^{(\ell)} + \alpha^{(\ell)} \mathbf{p}^{(\ell)} \\ &= \mathbf{x}^{(0)} + \sum_{i=0}^{\ell} \alpha^{(i)} \mathbf{p}^{(i)} \end{aligned}$$

Initially, one has: $\mathbf{p}^{(0)} = \mathbf{r}^{(0)} \Rightarrow \mathbf{p}^{(0)} \in \mathcal{K}_1(\mathbf{A}, \mathbf{r}^{(0)})$.

At iteration ℓ , one has: $\mathbf{p}^{(\ell+1)} = \mathbf{r}^{(\ell+1)} + \beta^{(\ell)} \mathbf{p}^{(\ell)}$

$$\begin{aligned} &= \mathbf{r}^{(\ell)} - \alpha^{(\ell)} \mathbf{A} \mathbf{p}^{(\ell)} + \beta^{(\ell)} \mathbf{p}^{(\ell)} \\ &= \mathbf{r}^{(0)} - \sum_{i=0}^{\ell} \alpha^{(i)} \mathbf{A} \mathbf{p}^{(i)} + \beta^{(\ell)} \mathbf{p}^{(\ell)} \end{aligned}$$

If $\mathbf{p}^{(i)} \in \mathcal{K}_{i+1}(\mathbf{A}, \mathbf{r}^{(0)})$ for $\forall i < \ell$, then $\mathbf{p}^{(\ell+1)} \in \mathcal{K}_{\ell+2}(\mathbf{A}, \mathbf{r}^{(0)})$.

Then:

$$\mathbf{x}^{(\ell+1)} - \mathbf{x}^{(0)} = \sum_{i=0}^{\ell} \alpha^{(i)} \mathbf{p}^{(i)} \in \mathcal{K}_{\ell+1}(\mathbf{A}, \mathbf{r}^{(0)})$$

□

Properties of the CG method (*continuation*)

$$\text{Property: } \mathbf{x}^{(\ell)} - \mathbf{x}^{(0)} \in \mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})$$

$$\text{Property: } \mathbf{x}^{(\ell)} = \mathbf{x}^{(0)} + \underset{\mathbf{y} \in \mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})}{\operatorname{argmin}} J(\mathbf{x}^{(0)} + \mathbf{y})$$

At each iteration ℓ , one has the best solution $\mathbf{x}^{(\ell)}$ in the sense “ J is minimum” such that $\mathbf{x}^{(\ell)} - \mathbf{x}^{(0)}$ belongs to the subspace $\mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})$.

Unfortunately, the CG method is limited to SPD/HPD matrices.

Toward general iterative methods for general matrices ...

We seek for a method that gives ...

$$\mathbf{x}^{(\ell)} - \mathbf{x}^{(0)} = \mathcal{P}_{\ell-1}(\mathbf{A}) \mathbf{r}^{(0)} \quad \text{where } \mathcal{P}_{\ell-1}(\cdot) \text{ is a polynomial of degree } \ell - 1$$

such that $\mathbf{x}^{(\ell)}$ is the “best solution” with $\mathbf{x}^{(\ell)} - \mathbf{x}^{(0)} \in \mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})$.

The **GMRES** (*generalized minimal residual*) method is a Krylov method based on the minimization of the residual at each iteration. (*There are other Krylov methods.*)

Nonstationary iterative methods for linear systems

Conjugate gradient method

Interlude on Krylov spaces

Few words about GMRES

GMRES — Principle

The **GMRES** (*Generalized Minimal Residual*) method relies on 2 steps performed at each iteration ℓ :

1. Building an orthonormal basis $\{\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(\ell)}\}$ for the Krylov subspace $\mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})$

$$\mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)}) := \text{span}(\mathbf{r}^{(0)}, \mathbf{A}\mathbf{r}^{(0)}, \dots, \mathbf{A}^{\ell-1}\mathbf{r}^{(0)}) = \text{span}(\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(\ell)})$$

With GMRES:

- The basis vectors are built by using Arnoldi iterations.
- Only the additional vector $\mathbf{v}^{(\ell)}$ is computed at iteration ℓ .

2. Solving a minimization problem to get the “best solution”

$$\mathbf{x}^{(\ell)} = \mathbf{x}^{(0)} + \underset{\mathbf{y} \in \mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})}{\text{argmin}} \|\mathbf{b} - \mathbf{A}(\mathbf{x}^{(0)} + \mathbf{y})\|_2$$

The solution $\mathbf{x}^{(\ell)}$ is such that:

- The update belongs to the Krylov subspace: $\mathbf{x}^{(\ell)} - \mathbf{x}^{(0)} \in \mathcal{K}_\ell(\mathbf{A}, \mathbf{r}^{(0)})$
- The 2-norm of the residual is minimum: $\|\mathbf{r}^{(\ell)}\|_2$ is minimum

With GMRES:

- Solving this problem is equivalent to solving a least square problem.
- The least square problem can be solved with a QR factorization.
- The QR factorization can be computed rapidly thanks to Givens matrices.

GMRES algorithm (*main steps*)

```

 $\mathbf{x}^{(0)} \in \mathbb{R}^n$ 
 $\mathbf{r}^{(0)} = \mathbf{b} - \mathbf{A}\mathbf{x}^{(0)}$ 
 $\mathbf{v}^{(1)} = \mathbf{r}^{(0)} / \|\mathbf{r}^{(0)}\|$ 
for  $\ell = 1, 2 \dots$  do
    // Building the orthonormal basis
     $\mathbf{w}^{(\ell)} = \mathbf{A}\mathbf{v}^{(\ell)}$ 
    for  $i = 1, \dots, \ell$  do
        |  $\mathbf{w}^{(\ell)} = \mathbf{w}^{(\ell)} - (\mathbf{w}^{(\ell)}, \mathbf{v}^{(i)})\mathbf{v}^{(i)}$ 
    end
    If  $\|\mathbf{w}^{(\ell)}\|_2 = 0 \rightarrow$  Stop.
     $\mathbf{v}^{(\ell+1)} = \mathbf{w}^{(\ell)} / \|\mathbf{w}^{(\ell)}\|_2$ 
    // Solving the minimization problem
     $\mathbf{z}^{(\ell)} = \operatorname{argmin}_{\mathbf{z} \in \mathbb{R}^{\ell+1}} \|\mathbf{b} - \mathbf{A}\mathbf{V}^{(\ell)}\mathbf{z}\|_2$ 
     $\mathbf{x}^{(\ell)} = \mathbf{x}^{(0)} + \mathbf{V}^{(\ell)}\mathbf{z}^{(\ell)}$ 
end

```

$\mathbf{V}^{(\ell)}$ is a $\ell \times (\ell + 1)$ matrix which the columns are the basis vectors $\{\mathbf{v}^{(i)}\}_{i=1 \dots \ell+1}$.

Theoretical aspects

- ▶ Method for general nonsingular matrices
- ▶ By construction, **convergence with maximum n iterations.** (*with ∞ accuracy*)
- ▶ If less iterations are required, procedure stopped during the construction of the basis. → **Breakdown**

Algorithmic aspects

- ▶ **The computational cost increases $\mathcal{O}(\ell^2)$ at each iteration.**
 - Storage of an additional basis vector and larger matrices
 - Orthogonalization by an additional basis vector
 - Solution of a larger minimization problem

To reduce the cost, the process can be restarted by using the current solution as initial solution. → **Restarted GMRES**

- ▶ **The algorithm is easy to parallelize.**
 - Basic linear algebra operations (BLAS 1 et 2) \Rightarrow Easy for parallel computing
 - Computation of scalar product and norms \Rightarrow Collective communications

Standard approach for nonsymmetric matrices. Widely used!
*Need to limit the number of iterations \Rightarrow **Preconditioning** ...*

Summary

- ▶ **Stationary methods** ($\mathbf{M}\mathbf{x}^{(k+1)} = \mathbf{N}\mathbf{x}^{(k)} + \mathbf{b}$)
 - **Jacobi** and **Gauss-Seidel (G.-S.)** methods
 - Improvements: “relaxation” (*JOR* and *SOR*) and “by block” approaches
 - Algorithmic aspects:
 - Matrix-vector products and linear combinations
 - Parallelism easy for Jacobi, a bit more complicated for G.-S.
 - Finite difference problem → *red/black approach* for G.-S.
- ▶ **Unstationary methods** ($\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha^{(k)}\mathbf{p}^{(k)}$)
 - **Steepest descent** and **Conjugate Gradient (CG)** methods
 - If \mathbf{A} SDP: link with quadratic optimisation, conv. in max. n iterations
 - Algorithmic aspects: matrix-vector prod., lin. comb., scalar products
- ▶ **Krylov methods** ($\mathbf{x}^{(k)} = \mathbf{x}^{(0)} + \mathbf{V}^{(k)}\mathbf{z}^{(k)}$) which are unstationary methods
 - **GMRES**
 - For nonsingular \mathbf{A} : conv. in max. n iterations
 - Algorithmic aspects: iterations of increasing cost, parallelism is easy

Resources

- ▶ *Numerical Linear Algebra*
L.N. Trefethen, D. Bau III (1997), SIAM
- ▶ *Iterative Methods for Sparse Linear Systems, 2nd edition*
Y. Saad (2003), SIAM
- ▶ *Méthodes Numériques : Algorithmes, analyse et applications*
A. Quarteroni, R. Sacco, F. Saleri (2007), Springer
- ▶ *Calcul scientifique parallèle*
F. Magoulès et F.-X. Roux (2017), Dunod
- ▶ *Calcul scientifique parallèle*
P. Ciarlet et E. Jamelot, polycopié de cours
- ▶ M. H. Gutknecht. “A Brief Introduction to Krylov Space Methods for Solving Linear Systems”, Proc. of the Int. Symp. on Front. of Comput. Sci. (2005) [[Preprint](#)]